



How to Sample Hay and Baleage

Test each lot of forage separately. A lot is defined as hay that was managed the same and harvested at the same time, out of the same field, and under the same conditions.

A forage analysis is only as good as the sample collected. For best results, collect a representative sample by following these guidelines:

Dry Hay



- Using a hay coring probe, collect 15–20 cores from each lot.
- To collect a core, insert the hay probe on the side of the bale, coring toward the center so you cross several windrows.
- Remove the hay probe and empty the contents into a clean container (e.g., bucket or bag).
- Mix the material thoroughly to decrease bias. Pour in enough sample to fill a 1-quart plastic bag.
- Label each sample to make it easy to remember the location and lot sampled.
- Fill out the [UGA Feed and Forage Testing Application](#), attach the form to the sample, and submit to your UGA County Extension Office.

Baleage

Baleage samples may be collected:

- **After baling, before wrapping:** Collect core samples just before wrapping, using the dry hay protocol.
- **After wrapping:** Collect core samples from various locations in one tube or several individually wrapped bales from the same lot. Ideally, this would occur post-fermentation (6+ weeks after wrapping).
- **Just before feeding:** Collect core samples from the tube or lot of bales that you plan to feed next. This decreases potential spoilage caused by hole punctures.

When sampling baleage:

-
- Seal the puncture hole in the bale immediately after coring with two pieces of UV-protected silage tape layered across each other to form an “X”.
 - Remove as much air as possible from the sample bag before sealing.
 - Ship baleage samples to the lab immediately. Collect and ship samples early in the week to reduce the chance of spoilage between collection and analysis.

How to Read a Forage Test Report

A forage test is one of the most valuable tools a producer has available. It is important to understand the terms on the report so you can interpret the results correctly.

As Sampled vs. Dry Matter: Which Do I Use?

A forage test will have two different numbers listed for each measure of forage quality. To compare numbers and ration balancing, use the numbers for dry matter.

Key Terminology from NIR Analysis

A forage test contains a lot of valuable information, but here are the main points of interest to producers.

Relative forage quality (RFQ): A calculation used to rank forages according to their relative nutritive value (see Table 1).

Table 1. Forage Quality Suggestions for Different Animal Classes.	Class	RFQ Range
	Heifer, 3–12 months	
	Heifer, 12–18 months	
	Heifer, 18–24 months	
	Dry cow	
	Stocker cattle	
	Lactating beef cow	
	Dairy calf	
	Dairy, first 200 days	
	Dairy, last 200 days	
	Idle horse	
	Brood mare	
	Nursing mare	
	Working horse	
	Hard-working horse	

Total digestible nutrients (TDN): A calculation of the energy value of the forage.

Neutral detergent fiber (NDF): A close estimate of the total fiber constituents of feedstuffs; this is negatively correlated to dry-matter intake.

Acid detergent fiber (ADF): Represents the least-digestible fiber portion and is used to calculate digestibility measures.

Acid detergent fiber (ADF): Represents the least-digestible fiber portion and is used to calculate digestibility measures.

Crude protein: Represents total nitrogen (including nitrates; see Table 2).

Table 2. Nitrates [†] in Animal Feed. Forage nitrate (ppm)	Strategy
< 4,500	safe to feed
4,500–6,500	limit to 1/2 ration for p
6,500–9,000	limit to 1/2 ration
9,000–15,000	limit to 1/3 ration
15,000–18,000	limit to 1/4 ration
> 18,000	potentially lethal, use

[†]Nitrates accumulate in forage with excessive nitrogen fertilization and stress that limits growth (e.g., moisture stress).

Forage Test Report Example

How to Use a Forage Test Report to Make a Winter Feed Plan for Beef Cattle Producers

Producers should follow these steps to develop a plan for winter feeding ahead of time and avoid trouble during the breeding season:

- Understand the cows' stage of production.
 - Cows should be rebred within 80–90 days postcalving to maintain a 365-day calving interval.
 - If the available forage does not meet a cow's basic nutrient requirements, she will breed later in the season. The basic requirements are outlined below.
- Understand the available forages. Ask the following questions:
 - How many bales of hay are available? How much does each bale weigh?
 - What is the nutritive value of the hay? Have you tested each lot of hay?
 - How long does the hay need to last?
- Develop an economical supplementation strategy to address nutrient deficiencies.
 - Evaluate available supplements carefully. They are not all created equally.
 - Evaluate the price of supplements on a dry-matter basis. Look at the cost per nutrient.
 - Utilize a program (e.g., [UGA Basic Balancer](#)) to accurately develop diets.

Table 3. Basic Nutrient Requirements of a 1,200 lb Cow.	Stage of producti on	DMI % of bw	CP %	TDN %
	Dry cow		1.86	
	Late gestation		1.86	
	Peak lactation		2.00	
	Late lactation		2.13	
<i>Note.</i> Required nutrients will increase or decrease as the cow's weight changes.				

Stage of production

DMI % of bw

CP %

TDN %

From "Nutrient Requirements of Beef Cattle, 8th
Press (www.doi.org/10.17226/19014).

Match Hay to Cow's Nutrient Requirements to Reduce Costs.

Reduce supplementation needs by feeding the best forage when the cow's requirements are the greatest. Numbers will change based on the feed source. Work with your local Extension agent to develop cows' diets.

Table 4. Example of Supplementation Needed Based on Forage Test Results.	Sta	Cu	Cu	Cu	
	ge	tti	tti	tti	
	of	ng	ng	ng	
	Pr	1	2	3	
	od	7%	9%	12	
	uct	CP	CP	%	
	ion	48	54	CP	
		%	%	60	
		TD	TD	%	
		N	N	TD	
				N	
				lb	
				supplement	
				required	
	Dry cow				0
	Late gestation				4
	Peak lactation				10
	Late lactation				6
	Note. Assuming 50:50 corn:hay				